



Donald W. Bennett, O.D., M.D.
Surgical and Consultative
Ophthalmology

B. Matthew Blair, M.D.
Surgical and Consultative
Ophthalmology

Steven M. Bloom, M.D., F.A.C.S.
Diseases and Surgery of the Retina
and Vitreous

Fraser McKay, O.D.
Consultative Optometry

Meredith Lanham, O.D.
Consultative Optometry

Ian W. McWherter,
O.D., F.A.A.O.
Consultative Optometry

John A. Millin, M.D.
Surgical and Consultative
Ophthalmology


Lee S. Peplinski, O.D., F.A.A.O.
Consultative Optometry

Inder P. Singal, M.D.
Diseases and Surgery of the Retina
and Vitreous

Andrew Steele, O.D.
Consultative Optometry

Lawrence R. Tenkman, M.D.
Corneal and Anterior
Segment Surgery

Nikolaos Zagorianos,
O.D., F.A.A.O.
Consultative Optometry



Amsler Grid

This test is designed to find abnormalities in the macula, the reading area of the eye. It helps to find many problems early when they may still be treatable.

Ideally the Amsler grid should be examined daily. It is often easier to remember to look at the grid if it is placed where you are sure to see it, such as on the refrigerator or the bathroom mirror.

Test each eye separately, covering the other eye. Glasses should be worn if they are normally needed for reading. You should look only at the central dot and while doing this, notice whether all of the lines are present. You should also look for any areas where the lines are wavy or distorted. If you notice any areas where the grid is missing, wavy, or distorted that are significantly different from your baseline, please call our office to schedule an examination within several days.

Toll Free to Louisville
1 (800) 477-0055
Fax (502) 361-4488

e-mail: admin@eyecenters.com
www.eyecenters.com